

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/337187183>

OROBANCHE CENTAURINA BERTOL. THE CORRECT NAME FOR O. KOCHII F.W. SCHULTZ (OROBANCHACEAE)

Article · November 2019

CITATION

1

READS

344

7 authors, including:



Óscar Sánchez Pedraja

Grupo botánico cantábrico (GBC), Liérganes, Spain

73 PUBLICATIONS 545 CITATIONS

[SEE PROFILE](#)



Luis Carlón Ruiz

52 PUBLICATIONS 513 CITATIONS

[SEE PROFILE](#)



Giannantonio Domina

Università degli Studi di Palermo

207 PUBLICATIONS 2,161 CITATIONS

[SEE PROFILE](#)



Renata Piwowarczyk

Jan Kochanowski University

123 PUBLICATIONS 876 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Index of Orobanchaceae [View project](#)



Phytotaxonomic investigation of the Mediterranean Area [View project](#)

OROBANCHE CENTAURINA BERTOL. THE CORRECT NAME FOR *O. KOCHII* F.W. SCHULTZ (*OROBANCHACEAE*)

**Jiří ZÁZVORKA¹, Óscar SÁNCHEZ PEDRAJA², Gonzalo MORENO MORAL³, Luis CARLÓN RUIZ⁴,
Gianniantonio DOMINA⁵, Manuel LAÍN Z GALLO⁶ & Renata PIWOWARCZYK⁷**

¹ Institute of Botany, Academy of Science of the Czech Republic. CZ-252 43 Průhonice (Czech Republic). zazvorka@ibot.cas.cz

² E-39722 Liérganes (Cantabria, Spain). osanchez@farmalierganes.com

³ C/Santa Clara, 9-1º dcha. E-39001 Santander (Cantabria, Spain)

⁴ Biosfera, Consultoría Medioambiental S. L. C/Candamo, 5. E-33012 Oviedo (Asturias, Spain)

⁵ Department of Agriculture, Food and Forest Sciences, University of Palermo, Viale delle Scienze, Bldg. 4. I-90128 Palermo (Italy). gianniantonio.domina@unipa.it

⁶ Avda. Hnos. Felgueroso, 25. E-33205 Gijón (Asturias, Spain). lainz@colegioinmaculada.es

⁷ Department of Botany, Institute of Biology, Jan Kochanowski University, 15 Świątokrzyska St. 25-406 Kielce (Poland). renata.piwowarczyk@ujk.edu.pl

ABSTRACT: After the studying the original material of *Orobanche centaurina* Bertol. (*Orobanchaceae*) deposited in BOLO, a forgotten species described from Massa (Central Italy) parasitizing *Centaurea paniculata* L., and compared it with the type of *O. kochii* F.W. Schultz and with the types of others taxa actually considered synonyms of this, we concluded that the name of Bertoloni is the correct one for this species. **Keywords:** Nomenclature; Eurasian flora; *Orobanchaceae*; parasitic plants.

RESUMEN: *Orobanche centaurina* Bertol., nombre correcto para *O. kochii* F.W. Schultz (*Orobanchaceae*). Después de estudiar el material original de *Orobanche centaurina* Bertol. (*Orobanchaceae*) depositado en BOLO, una olvidada especie descrita de Massa (Italia central) parásita de *Centaurea paniculata* L., y compararlo con el tipo de *O. kochii* F.W. Schultz y con los tipos de otros táxones actualmente considerados sinónimos de éste, estimamos que el nombre de Bertoloni es el correcto y prioritario para esta especie. **Palabras clave:** nomenclatura; flora euroasiática; *Orobanchaceae*; plantas parásitas.

INTRODUCTION

The renewed interest in the study of *Orobanche* and related genera, notoriously tricky plants, supported by the actual methodologies and equipments (*in vivo* colour photographs readily exchanged online and molecular tests for morphological hypotheses) have enabled a deeper reassessment of supposedly homogeneous species and the vindication of some previously unrecognised taxa. This happened with the broomrape taxa parasitizing *Centaurea* generically called *Orobanche major* L. or, after the formal rejection of this name as nomen ambiguum (TURLAND & RUMSEY, 1997), *O. elatior* Sutton (e.g. GRENIER, 1853; BECK, 1890; COSTE, 1904; BONNIER & DOUIN, 1926; BECK, 1930; NOVOPOKROVSKII & TZVELEV, 1958; CHATER & WEBB, 1972; TZVELEV, 1981; GILLI, 1982; TERYOKHIN & al., 1993; KREUTZ, 1995; UHLICH & al., 1995; ZHANG & TZVELEV, 1998; PUSCH, 2009; etc).

MATERIAL AND METHODS

A great number of herbarium specimens of the taxa under consideration were examined (ZÁZVORKA, 2010; PIWOWARCZYK & KRAJEWSKI, 2015; SÁNCHEZ PEDRAJA & al., 2016+) they are deposited in the following herbaria (B, BEOU, BOLO, BP, BRA, BRNM, BRNU, CB, CHRZ, CLF, CHOM, G, GM, GLM, KRA, KRAM, KTC, KTU, L, LBL, LE, LI, LINN, LOD, M, MPU, MW, P, PE, POZ, PR, PRC, ROZ, SAV, SLA, SLO, WA, WRSL, ZMT and

in private herbarium). Based on these herbarium specimens, observations of living plants and numerous images available on the Internet (see SÁNCHEZ PEDRAJA & al., 2016+), we have studied the taxonomy of the species parasites of *Centaurea* (s.l.) indiscriminately called *Orobanche major* L.

Broomrapes parasites on Centaurea in Europe and Africa, taxonomic story

In Spain, for instance, PUJADAS (1999) has reinstated *O. icterica* Pau, Not. Bot. Fl. España 3: 5. 1889 first at sub-specific rank (*O. major* subsp. *icterica* [Pau] A. Pujadas in Flora Montib. 11: 16. 1999) and soon after at the species level for the parasite of *Centaurea*, mainly *C. aspera*, traditionally referred to as *O. major*. His proposal gained wide acceptance, and most Spanish botanists nowadays recognize as a different species the *Centaurea*-parasite widespread in Spain, whose earliest name, as shown in CARLÓN & al. (2011), is the North-African *Centaurea*-parasite *O. leptantha* Pomel in Bull. Soc. Climatol. Alger 11: 110. 1874 [rosaceous form] (= *O. curvata* Pomel in Bull. Soc. Climatol. Alger 11: 111. 1874 [yellow form] (DOMINA & al., 2013; P 04363860!; P 03428431!)) another of those long buried taxa under the synonymy of *O. major* / *O. elatior*. (PUJADAS, 2013) contests this conclusion, but using the type specimen of *O. leptantha* as his only quantitative reference (we have seen other African materials readily contradicting the numbers in his table 1) and misrepresenting qualitative traits supposedly conflicting with Po-

mel's protologue and materials (for instance, the bracts of the Iberian plants characteristically exceed the corolla, and they can hardly be described as *ovate to triangular*).

But the Western Mediterranean region is far from being the only area in which the sequential all-purpose use of *O. major* and *O. elatior* proved to cover an artificial hodgepodge. The work by ZÁZVORKA (2010) showed beyond any doubt that two utterly different species (the true *O. elatior* Sutton in Trans. Linn. Soc. London 4: 178, t. 17. 1798 and *O. kochii* F.W. Schultz, Flora [Regensb.] 30(5): 66. 1847) had been hidden under a single name as a result of systematic host-driven misidentifications. Despite their similarity and even overlap in host specificity (*Centaurea scabiosa*, the main if not the only host of the former, is frequently attacked by the latter), to their morphological and molecular disparity they add a neat biogeographical divergence: *O. elatior* is a temperate-subatlantic species restricted to western, central and northern Europe (ZÁZVORKA, 2010; PIWOWARCZYK & KRAJEWSKI, 2015; SÁNCHEZ PEDRAJA & al., 2016+), whereas *O. kochii* is an Eurasian species which occurs in warmer summer regions, stretching from Mediterranean Europe (France and Italy) to perhaps China and India (ZÁZVORKA, 2010; PIWOWARCZYK & KRAJEWSKI, 2015; SÁNCHEZ PEDRAJA & al., 2016+).

In the present note we will keep pulling this thread in order to establish the correct nomenclature and summarize the geographic and host range of European *Centaurea* parasites.

In France we could personally observe that *O. kochii* does not only occur in the Alpine area with less Atlantic influence, but is widespread in sunny, dry, sandy spots across the strictly Mediterranean region, where as already stated by REICHENBACH f. (1862, sub *O. Ritro*, "Sur le *Centaurea aspera*, l'*Echinops Ritro*: sables de Mazarque [Marseille-Mazargues]. Roux et Blaise! [P 04419058!, P 04406071] ... Marseille in vineis. Sonder!); COSTE (1904, f. 2820 [*O. kochii* (see ZÁZVORKA, 2010, "straighter corolla back in the middle")], sub *O. major* L., p.p., "Sur les *Centaurea* et *Echinops*, dans l'Est, le Centre et le Midi [SLA 0484151!]) and BONNIER & DOUIN (1926, sub *O. major* L., p.p., "L'espèce est parasite sur les *Centaurea Scabiōsa*, *Centaurea aspera*, *Centaurea collina* ... Provence et Alpes-Maritimes, Languedoc") it parasitizes both *Echinops* and several species of *Centaurea*, namely *C. aspera* L.

Four taxa suspected to be conspecific with *O. kochii* are: *O. centaureae-scabiosae* F.W. Schultz., *O. centaurina* Bertol., *O. elatior* var. *forojuliensis* Coss. and *O. ritro* Gren.

Orobanche centaureae-scabiosae F.W. Schultz in Bot. Lit. Blätt. 5: 500. 1830, is based on *Orobanche de la Centaurée scabieuse* Vaucher, Monogr. Orobanch.: 61. 1827 and refers to a species described near Ronco Scrivia (Liguria) but according to Art. 36 of the ICN is not validly published because this name is listed as a mere synonym of *O. elatior*. Although the characters of the plant described by Vaucher do not coincide with those of a normally developed *O. centaurina*, we have made some attempts to know it. However, the meticulous efforts to locate the original material in G have been unsuccessful (L. Gautier 2017, pers. comm., 21 April) and our fieldworks dedicated to finding specimens of *Centaurea scabiosa* parasitized by broomrapes in the surroundings of Ronco Scrivia (Cipollina, Banchetta, Porale, Alpe di Porale,

Tana d'Orso), carried out on 4 July 2017, has not offered positive results.

Orobanche centaurina Bertol., Fl. Ital. 6: 430. 1846 (fig. 1), lectotype BOLO (DOMINA & MAZZOLA, 2011), is a taxon described from S. Giuseppe near Massa (Tuscany), on *C. paniculata* L. and in a similar thermophilous coastal grasslands in which *O. kochii* occurs in France.



Fig. 1. *O. centaurina* Bertol. The lectotype in BOLO designated by Domina & Mazzola (2011).

Orobanche elatior var. *forojuliensis* Coss., Notes Pl. Crit.: 8. 1849 (fig. 2); ≡ *O. forojuliensis* [(Coss.) Nyman] ex Beck in Engl., Pflanzenr., Orobanch.: 247. 1930, pro syn., lectotype (here designated) [the sheet contains two specimens, the right-hand specimen is a suitable lectotype for *O. elatior* var. *forojuliensis* Coss.]: 1. "E. Bourgeau, Env. de Fréjus. n.° 302, *Orobanche elatior*, Sutton ex Reut. in Prodr. var. *maritima*, Coss. ined., parasite sur le *Centaurea aspera*!, bords de la route entre Fréjus et S.^a Raphaël [Provence-Alpes-Côte d'Azur], 4 Juin, 1848" (P 04385331); isolectotypes (P 04384972 - Hb. E. Cosson [only the central specimen, close to its label]) and (P 04384973 - Hb. E. Cosson). All these materials support beyond any reasonable doubt that Cosson's trinomen is a mere synonym of *O. centaurina*.

Orobanche ritro Gren. in Gren. & Godr., Fl. France 2: 635. 1853 (fig. 3). Despite its vindication by PUJADAS (2013), a glance at the type (P 00654040!), and other sheets collected by Grenier in the same locality (near Gap, Provence-Alpes-Côte d'Azur), e. g. the topotype (P 0440 6079), moreover the illustrations of *O. ritro* Gren. in Gren. & Godr., Fl. France 2: 635. 1853 published by REICHENBACH f., Iconogr. Fl. Germ. Helv. 20: t. 170. 1862 and BONNIER, Fl. ill. France 8(78): t. 466 f. 2160b. 1926 (all showing ovate to triangular cauline leaves, corolla with straight dorsal line, divergent spoon-shaped lower lip lobes and flat, straight, subentire upper lip) and numerous photographs, in the nature, of the broomrape parasitizing both *Centaurea* and *Echinops* in the *terra classica* of *O. ritro*

(Provence-Alpes-Côte d'Azur), sent to us by Daniel Pavon, suffice to regard Grenier's binomial as a mere synonym of *O. kochii*, a claim otherwise supported by local botanists in recent times [ROYER & al., 2014; PAVON, 2015; PAVON & al., 2015; "around here (Bouches-du-Rhône) there's only one variable species" (D. Pavon 2017, pers. comm., 2 February)].




Fig. 2. *O. elatior* var. *forojuliensis* Coss.
The lectotype in P, here designated.

RESULTS

The Bertoloni's type refers to the same taxon of *O. kochii*, the two specimens of Bertoloni's sheet (fig. 1) show the typical characters of this species, stems relatively shorter with leaves broad at base and ovate-triangular shape, inflorescences subcylindrical, ± loose, corolla with straight dorsal line, lower lip with lobes divergent and spoon-shaped, upper lip flat, straight, subentire, specimens brownish-rusty when dry (see ZÁZVORKA, 2010: 82 t. 1). We could observe all these features even more unmistakably in a set of photographs of living plants taken in the Carrara marble quarries near the caves of Fantiscritti, about 10 km away from the *locus classicus* of *O. centaurina*, shared online in "Il Forum dei Funghi e Fiori in Italia" of the Associazione Micologica Italiana Naturalistica Telematica – AMINT [(https://www.funghiitaliani.it/), accessed: 21 Dec 2005, 23 Aug 2006] by the user "Apuano" and now removed. "Apuano" mentioned several species existing in the vicinity of the broomrape, the first was *Centaurea arachnoidea* Viv. (= *C. rupestris* L.). We'd be pleased to better acknowledge the

authorship of these pictures (fig. 4), but the user "Apuano" no longer belongs to the association and they have not address to contact him.



Fig. 3. *O. ritro* Gren. The lectotype in P, designated by Pujadas (2013).

The path would thus be paved for the name *O. centaurina* Bertol. to be adopted as the correct one for the plant so far called *O. kochii* (ZÁZVORKA, 2010; CARLÓN & al., 2011; PIWOWARCZYK, 2012; PIWOWARCZYK & KRAJEWSKI, 2015).

DISCUSSION

Summing up, three species of *Orobanche* can be found parasitizing *Centaurea* s.l. (incl. *Cyanus*, *Psephellus*, *Rhaponticoides*) in Europe: 1) *O. elatior* Sutton, with dense spikes of regularly curved corollas, divergent elliptic lower lip lobes and bracts shorter or slightly longer than the flowers, occurs to the northwest of an axis linking the Alps to the Pyrenees, where it is parasitic mainly if not only on *Centaurea scabiosa*; 2) *O. leptantha* Pomel (= *O. curvata* Pomel [described on a yellow variant] = *O. icterica* Pau), similar to *O. elatior* but with rounded, convergent lower lip lobes and longer bracts, often long exceeding the flowers, occurs around the western Mediterranean where it is parasitic on several species of *Centaurea* (mainly *C. aspera* L. but also *C. fragilis* Durieu, *C. hyssopifolia* Vahl, *C. ornata* Willd., *C. pullata* L. and *C. sphaerocephala* L.), as well as *Rhaponticoides alpina* (L.) M.V. Agab. & Greuter and *Cheirolophus intybaceus* (Lam.) Dostál; and 3) the distantly related *O. centaurina* Bertol. (= *O. kochii* F.W. Schultz = *O. ritro* Gren. [described on a yellow variant

common in this group], with dorsally straight flowers showing a flattened adaxial surface and big, divergent, spatulate, concave lower-lip lobes and patent, wide upper-lip lobes, is a “steppic” species occurring to the southeast of the above-mentioned imaginary axis, where it parasitizes chiefly *Echinops* and *Centaurea* (mainly *C. scabiosa* and *C. paniculata* s.l.) and apparently also *Rhaponticoides ruthenica* (Lam.) M.V. Agab. & Greuter (PLANTARIUM, 2007+, Russian republic of Tatarstan) and *Ptilostemon echi-nocephalus* (Willd.) Greuter (PLANTARIUM, 2007+, Crimean Peninsula).



Fig. 4. *Orobanche centaurina* Bertol. Left photo: near Fantiscritti (10 km north of Massa, *loc. class.*), Italy, probably parasite of *Centaurea rupestris*, “Apuano”, 2005 (sub *O. elatior*). Right photo: military maneuvers camp of Aglanet, pr. Orange (Provence-Alpes-Côte d’Azur, *terra class.* of *O. elatior* var. *forjuliensis*), France, 31TFJ4587, 50 m, beside *Centaurea aspera* in very dry and sunny soils, L. Carlón, G. Gómez, M. Laínz, G. Moreno MM0106/2003 & J.-M. Tison, 28 May 2003 (hb. Sánchez Pedraja 11371). Photos “Apuano” and G. Moreno.

Acknowledgements: We would like to express our gratitude to all those who kindly put their herbarium material at the authors’ disposal. Our thanks are especially due to the curators of herbaria BOLO, G, P and SLA for their assistance.

REFERENCES

- BECK, G. (1890) *Monographie der Gattung Orobanche*. Biblioth. Bot. 19. Cassel: Theodor Fischer.
- BECK, G. (1930) Orobanchaceae. In ENGLER, A. (ed.) *Das Pflanzenreich. Regni vegetabili conspectus* 96(IV.261): 1-348. Leipzig: Wilhelm Engelmann.
- BONNIER, G.E.M. & DOUIN, R.C.V. (1926) *Flore Complète illustrée en couleurs de France, Suisse et Belgique (comprenant la plupart des plantes d’Europe)* 8(78): 89-102, t. 463-467. Neuchatel: Delachaux et Niestlé; Paris: Librairie Générale de l’Enseignement, E. Orlhac.
- CARLÓN, L., LAÍNZ, M., MORENO MORAL, G. & SÁNCHEZ PEDRAJA, Ó. (2011) A new species (*Orobanche loscosii*), a priority name for *O. icterica* (*O. leptantha*) and a new member of the spanish flora (*O. elatior*). *Fl. Montib.* 48: 89-101.
- CHATER, A.O. & WEBB, D.A. (1972) *Orobanche* L. In TUTIN, T.G. & al. (eds.) *Flora Europaea* 3: 286-293. Cambridge University Press.
- COSTE, H. (1904) *Flore descriptive illustrée de la France, de la Corse et des contrées limitrophes* 3(1): 1-96. Paris: Librairie des Sciences naturelles Paul Klincksieck.
- DOMINA, G. & MAZZOLA, P. (2011) Notes on the genus *Orobanche* in Italy: 3. Taxa described by A. Bertoloni. *Plant. Biosystems* 145(2): 342-346.
- DOMINA, G., GREUTER, W., MARINO, P. & SCHÄFER, P. A. (2013) Types of names of *Orobanche* taxa described from North Africa. *Plant Biosystems* 147(3): 758-766.
- GBIF (The Global Biodiversity Information Facility). GBIF Online Resource Centre. Orobanchaceae. <http://www.gbif.org/species/7332526> [accessed: Jan 2017].
- GILLI, A. (1982) *Orobanchaceae*. In Davis, P.H. (ed.) *Flora of Turkey and the East Aegean Islands* 7: 1-23. Edinburgh: Edinburgh University Press.
- GRENIER, J.C.M. (1853) Orobanchées. In GRENIER, J.C.M. & GODRON, D.A. (1847-1856) *Flore de France, ou description des plantes qui croissent naturellement en France et en Corse* 2(2): 623-624. Paris: J.-B. Baillière.
- KREUTZ, C.A.J. (1995) *Orobanche: die Sommerwurzarten Europas: ein Bestimmungsbuch [The European broomrape species: a field guide]*. Vol. 1 [Mittel- und Nordeuropa (Central and Northern Europe)]. Maastricht: Stichting Natuurpublicaties Limburg.
- MNHN (Muséum National d’Histoire Naturelle de Paris) (2017) Botanique. Plantes vasculaires. <http://www.mnhn.fr/fr/collections/ensembles-collections/botanique/plantes-vasculaires> [accessed: Jan 2017].
- NOVOPOKROVSKII, I.V. & TZVELEV, N.N. (1958) *Orobanchaceae* Vent. In KOMAROV, V.L. (ed.) 1934-1964. *Flora SSSR* 23: 18-127. Moskva-Leningrad: Akad. Nauk SSSR.
- PAVON, D. (2015) Contribution à la connaissance et à la conservation des orobanches du département des Bouches-du-Rhône. *Bull. Soc. linn. Provence* 66: 57-88.
- PAVON, D., MICHAUD, H., VELA, E. & TISON, J.M. (2015) *Orobanche staehelinae* (Orobanchaceae), a new species from southeast France. *Phytotaxa* 207(1): 93-105.
- PIWOWARCZYK, R. (2012) The genus *Orobanche* L. (Orobanchaceae) in the Małopolska Upland (S Poland): distribution, habitat, host preferences, and taxonomic problems. *Biodiv. Res. Conserv.* 26: 3-22.
- PIWOWARCZYK, R. & KRAJEWSKI, Ł. (2015) *Orobanche elatior* and *O. kochii* (Orobanchaceae) in Poland: distribution, taxonomy, plant communities and seed micromorphology. *Acta Soc. Bot. Poloniae* 84: 103-123.
- PLANTARIUM (2007+) *Открытый атлас сосудистых растений России и сопредельных стран [Open Atlas of Vascular Plants of Russia and Adjacent Countries]*. *Определитель растений on-line [The determinant of plants on-line]*. *Scrophulariaceae. Orobanche*. <http://www.plantarium.ru/page/taxonomy/taxon/44566.html> [accessed: Jan 2017].
- PUJADAS, A. (1999) *Orobanche icterica* Pau, taxon minusvalorado del Sistema Ibérico. *Fl. Montib.* 11: 15-18.
- PUJADAS, A. (2013) *Orobanche icterica* Pau and *Orobanche ritro* Gren. & Godr. (Orobanchaceae) in the Iberian Flora. *Acta Bot. Malacitana* 38: 160-162.
- REICHENBACH, H.G.L. (1834-1914) *Icones florum germanicae et helveticae ...* 20: 49-125, pl. 121-220 (1862) [H.G. REICHENBACH f.]. Lipsiae: sumptibus Ambrosi Abel.
- ROYER, J.-M., TISON, J.-M. & MISSET, C. (2014). *Orobanche*. In TISON, J.-M. & FOUCAULT, B. de (2014) *Flora Gallica. Flore de France*. Biotope Éditions.
- SÁNCHEZ PEDRAJA, Ó., MORENO MORAL, G., CARLÓN, L., PIWOWARCZYK, R., LAÍNZ, M. & SCHNEEWEISS, G.M. (2016+) *Index of Orobanchaceae*. <http://www.farmalierganes.com/Otrospdf/publica/Orobanchaceae%20Index.htm>. [accessed: Jan 2017].
- TERYOKHIN, E.S., SHIBAKINA, G.V., SERAFIMOVICH, N.B. & KRAVTSOVA, T.I. (1993). In BUDANTSEV, L.Y.

- (ed.) *Opredelitel zarazikhovykh flory SSSR (Determinator of Broomrapes of the USSR Flora)*. Leningrad: Nauka Publ.
- TURLAND, N. & RUMSEY, F. (1997) (1318-1319) Proposals to reject the names *Orobanche major* and *O. laevis* (*Orobanchaceae*). *Taxon* 46(4): 787-791.
- TZVELEV, N.N. (1981) *Orobanchaceae* Vent. In FEDOROV An.A. (ed.). *Flora of Russia: The European Part and Bordering Regions* 5: 434-461. Leningrad: Nauka Publ.
- UHLICH, H., PUSCH, J. & BARTHEL, K.-J. (1995) *Die Sommerwurzen Europas: Gattung Orobanche*. Westarp Wissenschaften. Magdeburg.
- ZÁZVORKA, J. (2010) *Orobanche kochii* and *O. elatior* (*Orobanchaceae*) in central Europe. *Acta Musei Moraviae, Sci. Biol. (Brno)* 95(2): 77-119.
- ZHANG, Z.Y. & TZVELEV, N.N. (1998) *Orobanchaceae* Ventenat. In WU, Z.Y. & RAVEN, P.H. (eds.). *Flora of China* 18: 229-243. <http://www.efloras.org/> [accessed: Jan 2017].



Fig. 5. *Orobanche centaurina* Bertol. Boria, near Ćmielów (Małopolska Upland, Poland, original area of distribution of *O. kochii*), on *Centaurea scabiosa*, xerothermic grassland, R. Piwowarczyk, 1 July 2014 (hb. KTC). Photo R. Piwowarczyk.




Fig. 6. *Orobanche centaurina* Bertol. Boria, near Ćmielów (Małopolska Upland, Poland; original area of distribution of *O. kochii*), on *Centaurea scabiosa*, xerothermic grassland, R. Piwowarczyk, 1 July 2014 (hb. KTC). Photo R. Piwowarczyk.



Fig. 7. *Orobanche centaurina* Bertol. Near Quartier du Grès, south of Orange (Provence-Alpes-Côte d'Azur, *terra class.* of *O. elatior* var. *forojuliensis*), France, 31TFJ4584, 55 m, beside *C. aspera* in roadside ditch, G. Gómez & G. Moreno MM0121/2006 (hb. Sánchez Pedraja 12593). Photo G. Moreno.

(Recibido el 1-VIII-2019)
(Aceptado el 2-IX-2019)



Guía imprescindible de las flores de la Depresión del Ebro

Javier Puente Cabeza

Col. *Guías imprescindibles de flora*, nº 5
Encuadernación rústica cosida 11 × 21,6 cm
380 páginas en **COLOR**

Fecha lanzamiento: **julio de 2018**

ISBN: 978-84-947985-3-5

PVP: 24,00€ + envío

Estudio monográfico sobre los géneros *Hieracium* y *Pilosella* en España

Con referencias a Portugal y los Pirineos franceses

Gonzalo Mateo y Fermín del Egado

Monografías de Botánica Ibérica, nº 20


Encuadernación rústica cosida 17 × 24 cm

422 páginas en B/N y **COLOR**

Fecha lanzamiento: **enero de 2018**

ISBN: 978-84-945880-8-2

PVP: 30€- + envío



Flora vascular del término municipal de Córdoba *Catálogo florístico y claves de identificación*


Javier López Tirado

Monografías de Botánica Ibérica, nº 2
Encuadernación rústica cosida 17 × 24 cm
374 páginas en B/N y color

Fecha lanzamiento: **abril de 2018**

ISBN: 978-84-947985-0-4

PVP: 22,50€ + envío



Orchids of Central Spain (Cuenca Province). A Field Guide

Agustín Coronado & Eduardo Soto

Col. Essential Guides of Flora, nº 2

Encuadernación rústica 14,8 × 21 cm

244 páginas en **COLOR**

Fecha lanzamiento: **marzo de 2019**

ISBN: 978-84-947985-2-8

PVP: 25,95€ + envío

Mapa de vegetación actual, a escala 1: 10.000, del Parque Nacional de Ordesa y Monte Perdido y su zona periférica de protección

José Luis BENITO ALONSO



Organismo Autónomo Parque Nacionales

Encuadernación cartoné cosida 17 × 24 cm

450 páginas en **COLOR**

Fecha lanzamiento: **diciembre de 2018**

ISBN: 978-84-8014-916-7



Topónimos y apellidos españoles de origen ibérico o pre-latino

(Los iberos seguimos aquí)

Gonzalo Mateo Sanz

Monografías de Toponimia Ibérica, nº 1

Encuadernación rústica cosida 17 × 24 cm

230 páginas en **B/N**

Fecha lanzamiento: **junio de 2019**

ISBN: 978-84-947985-9-7

PVP: 15€ + envío

Manual para el trabajo de campo del proyecto GLORIA

Aproximación al estudio de las cimas. Métodos básico, complementarios y adicionales. 5ª edición

Harald Pauli & al.


Encuadernación rústica A4

150 páginas en **COLOR**

Fecha lanzamiento: abril de 2019

ISBN: 978-84-947985-7-3

PVP: 15€ + envío



Robles, hombres y dioses

Usos y visión simbólica de las quercus en Europa

Jesús IZCO

Monografías de Botánica Ibérica, nº 19

Encuadernación rústica cosida 17 × 21,9 cm

424 páginas en color

Fecha lanzamiento: febrero de 2016

ISBN: 978-84-945880-3-7

PVP: 29,95€ + envío

Flora vascular del Parc Natural del Túrria

Aurelio Peña, P. Pablo Ferrer, Jesús Riera, Javier Fabado & Gonzalo Mateo

Encuadernación rústica cosida 17 × 24 cm


249 páginas en **COLOR**

Idioma: valenciano

Fecha lanzamiento: octubre de 2017

ISBN: 978-84-945880-4-4

PVP: 25,00€ + envío



Orquídeas de Aragón

Conchita MUÑOZ ORTEGA

Col. *Guías imprescindibles de flora*, nº 2



Encuadernación rústica cosida 10 x 21 cm

202 páginas **en color con 250 fotografías**

Primera edición: abril de 2014

ISBN: 978-84-941996-1-5

PVP: 17,50 € + envío



Guía imprescindible de las flores del Prepirineo

Javier PUENTE CABEZA & José Luis BENITO ALONSO

Col. *Guías imprescindibles de flora*, nº 3

Encuadernación rústica cosida 17 x 24 cm

204 páginas **en color con más de 530 fotografías.**

Primera edición: abril de 2013

ISBN: 978-84-941996-4-6

PVP: 17,50 € + envío

Orquídeas de la provincia de Cuenca

Guía de campo

Agustín Coronado Martínez y Eduardo Soto Pérez

Colección Guías imprescindibles de flora, 4

Encuadernación rústica cosida 14,8 x 21 cm

252 páginas **en COLOR**

Fecha lanzamiento: mayo de 2017

ISBN: 978-84-945880-5-1

PVP: 25,95€ + envío

